

Using self-report surveys at the beginning of service to develop multi-outcome risk models for new soldiers in the U.S. Army

A. J. Rosellini¹, M. B. Stein^{2,3}, D. M. Benedek⁴, P. D. Bliese⁵, W. T. Chiu¹, I. Hwang¹, J. Monahan⁶, M. K. Nock⁷, M. V. Petukhova¹, N. A. Sampson¹, A. E. Street^{8,9}, A. M. Zaslavsky¹, R. J. Ursano⁴, R.C. Kessler^{1*} and On behalf of the Army STARRS Collaborators

¹Department of Health Care Policy, Harvard Medical School, Boston, Massachusetts, USA

²Departments of Psychiatry and Family Medicine & Public Health, University of California San Diego, La Jolla, California, USA

³VA San Diego Healthcare System, San Diego, CA, USA

⁴Department of Psychiatry, Center for the Study of Traumatic Stress, Uniformed Services University School of Medicine, Bethesda, MD, USA

⁵Darla Moore School of Business, University of South Carolina, Columbia, South Carolina, USA

⁶School of Law, University of Virginia, Charlottesville, VA, USA

⁷Department of Psychology, Harvard University, Cambridge, Massachusetts, USA

⁸National Center for PTSD, VA Boston Healthcare System, Boston, Massachusetts, USA

⁹Department of Psychiatry, Boston University School of Medicine, Boston, Massachusetts, USA

Background. The U.S. Army uses universal preventives interventions for several negative outcomes (e.g. suicide, violence, sexual assault) with especially high risks in the early years of service. More intensive interventions exist, but would be cost-effective only if targeted at high-risk soldiers. We report results of efforts to develop models for such targeting from self-report surveys administered at the beginning of Army service.

Methods. 21 832 new soldiers completed a self-administered questionnaire (SAQ) in 2011–2012 and consented to link administrative data to SAQ responses. Penalized regression models were developed for 12 administratively-recorded outcomes occurring by December 2013: suicide attempt, mental hospitalization, positive drug test, traumatic brain injury (TBI), other severe injury, several types of violence perpetration and victimization, demotion, and attrition.

Results. The best-performing models were for TBI (AUC=0.80), major physical violence perpetration (AUC=0.78), sexual assault perpetration (AUC=0.78), and suicide attempt (AUC=0.74). Although predicted risk scores were significantly correlated across outcomes, prediction was not improved by including risk scores for other outcomes in models. Of particular note: 40.5% of suicide attempts occurred among the 10% of new soldiers with highest predicted risk, 57.2% of male sexual assault perpetrations among the 15% with highest predicted risk, and 35.5% of female sexual assault victimizations among the 10% with highest predicted risk.

Conclusions. Data collected at the beginning of service in self-report surveys could be used to develop risk models that define small proportions of new soldiers accounting for high proportions of negative outcomes over the first few years of service.

Received 10 February 2017; Revised 28 February 2017; Accepted 28 February 2017

Key words: Army, disciplinary problems, mental health, military, predictive modeling, risk assessment, violence.

Introduction

The U.S. Army and Department of Defense (DoD) have implemented prevention programs for several negative outcomes (Department of the Army, 2015b), including suicide (Department of the Army, 2015a), workplace

violence (Department of Defense, 2014b), and sexual assault (Department of Defense, 2014a), that have high prevalence in the early years of the Army career (Kaufman *et al.* 2000; Department of the US Army, 2010, 2012). These preventive interventions are mostly universal; that is, all personnel are required to participate and the interventions are relatively non-intensive. More intensive interventions exist and could be implemented (Vungkhanching *et al.* 2007; Parkkari *et al.* 2011; Shea *et al.* 2013; Rudd *et al.* 2015; Senn *et al.* 2015), but would be cost-effective only if targeted at high-risk

* Address for correspondence: R. C. Kessler, Ph.D., Department of Health Care Policy, Harvard Medical School, 180 Longwood Avenue, Boston, MA, USA.
(Email: Kessler@hcp.med.harvard.edu)

personnel (Foster & Jones, 2006; Golubnitschaja & Costigliola, 2012). This targeting would require valid risk prediction tools. Recent studies have shown that Army and DoD administrative data can be used to develop such tools to predict negative soldier outcomes such as suicide (Kessler *et al.* 2015), violent crime perpetration (Rosellini *et al.* 2016), and sexual assault victimization (Street *et al.* 2016), but these models are limited by the fact that administrative data only become available over the course of time and are unavailable when preventive interventions might most logically be implemented at the beginning of service. An alternative would be to implement a risk factor survey at the beginning of service to target new recruits for preventive interventions. The current report presents the results of an attempt to develop risk models for a number of high-priority negative outcomes using data collected in such a survey of new U.S. Army soldiers subsequently followed over the first 2 years of service. If successful, the logic of this approach might be generalizable to a wide range of other workplace settings.

Methods

Sample

The survey was the New Soldier Survey (NSS) of the *Army Study to Assess Risk and Resilience in Servicemembers* (Army STARRS) (Ursano *et al.* 2014). The NSS was implemented April 2011–November 2012 in a representative sample of new U.S. Army soldiers prior to beginning Basic Combat Training (BCT) at Fort Benning, GA, Fort Jackson, SC, and Fort Leonard Wood, MO. Recruitment began by selecting weekly samples of 200–300 new soldiers at each installation to attend an informed consent presentation within 48 h of reporting for BCT. The presentation explained study purposes, confidentiality, voluntary participation, and answered all attendee questions before seeking written informed consent for a self-administered computerized questionnaire (SAQ) and neurocognitive tests and to link these data to the soldier's administrative records. These study recruitment and consent procedures were approved by the Human Subjects Committees of all Army STARRS collaborating organizations. The 21 832 NSS respondents considered here represent all Regular Army soldiers who completed the SAQ and agreed to administrative data linkage (77.1% response rate) (Rosellini *et al.* 2015). Data were doubly-weighted to adjust for differences in survey responses among the respondents who did *v.* did not agree to administrative record linkage and differences in administrative data profiles between

the latter subsample and the population of all new soldiers.

Outcomes

Outcome data were abstracted from 14 administrative databases through December 2013 (13–33 follow-up months after NSS completion) to operationalize 12 high-priority outcomes involving mental–physical disorders (Canham-Chervak *et al.* 2010; Department of the US Army, 2010; Institute of Medicine, 2010), violent crime perpetration–victimization (Institute of Medicine, 2010; Department of the US Army, 2012), and career problems (Kubisiak *et al.* 2009; Kapp, 2013). Dichotomous dependent variables were defined for first occurrence of each of the following outcomes:

Mental and physical disorders

Suicide attempt was defined based on the DoD Suicide Event Reporting system and International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) treatment codes. A positive test for illicit drug use was defined based on the Army substance abuse tracking database. Three health outcomes were defined from electronic medical records: mental hospitalization, traumatic brain injury (TBI); and any other severe injury exclusive of TBI (e.g. blindness, deafness, amputation, severe burns, paralysis), all based on ICD-9-CM codes.

Violent crime

DoD criminal justice databases were used to define three measures of violent crime perpetration [major physical, sexual assault, and minor violence (e.g. harassment)] and two of victimization (minor violence, sexual assault) coded according to the Bureau of Justice Statistics National Corrections Reporting Program classification system (U.S. Department of Justice, 2011) and of sufficient frequency to develop prediction models. The perpetration outcomes were defined from records of 'founded' offenses (i.e. where the Army found sufficient evidence to warrant full investigation). The victimization outcomes were defined using *any* officially reported victimization regardless of evidence.

Army career problems

Premature attrition from service due to career or personal problems (e.g. a character or behavior disorder, disability) was defined using an Army personnel database that tracks reasons for separating from service. Demotion was defined using information in the Army master personnel file.

Independent variables

Our goal was to optimize classification of soldiers who subsequently had the outcomes rather than to test specific hypotheses. As a result, we considered all potential predictors for which there was any evidence in the literature; e.g. risk-protective factors for suicidality (Nock *et al.* 2013; Afifi *et al.* 2016), mental hospitalization (Iribarren *et al.* 2000; Ryttila-Manninen *et al.* 2014), substance problems (Kilpatrick *et al.* 2000; Kirst *et al.* 2014), TBI (Cassidy *et al.* 2004; Elmasry *et al.* 2017), other severe physical injuries (Bulzacchelli *et al.* 2014; Theodoroff *et al.* 2015), violence perpetration (Dahlberg, 1998; Elbogen *et al.* 2010), violence victimization (Suris & Lind, 2008; Turchik & Wilson, 2010), and career problems (Knapik *et al.* 2004; Booth-Kewley *et al.* 2010). In total, 727 independent variables were operationalized from the SAQ in addition to eight performance-based neurocognitive test measures assessed in conjunction with the SAQ and 37 basic administrative variables recorded for all new soldiers at the beginning of service (772 total variables) (online Supplementary Appendix Table 1 summarizes all independent variables).

The SAQ variables were in six categories that included: sociodemographics (e.g. age, sex, race-ethnicity), self-reported lifetime history of DSM-IV mental disorders (ADHD, bipolar disorder, conduct disorder, generalized anxiety disorder, major depressive disorder, oppositional-defiant disorder, panic disorder, PTSD, substance abuse-dependence), exposure to stressors (childhood adversities, other lifetime traumatic stressors, past-year stressful life events and difficulties), personality (e.g. neuroticism, impulsivity, secure attachment), social networks (e.g. number of friends, number of sexual partners), and lifetime suicidality/non-suicidal self-injury (referred to henceforth as 'self-harm').

The neurocognitive variables, described in more detail elsewhere (Moore *et al.* 2017), assessed seven constructs: mental flexibility, attention, working memory, impulse control, facial memory, emotion identification, and bias toward negative emotions. A standardized efficiency score (the average of test accuracy and speed) was defined for each neurocognitive construct along with a composite overall efficiency score across the seven constructs. The administrative variables included Armed Forces Qualification Test (AFQT) scores, physical profile system (PULHES) scores, enlistment military occupational specialty classifications, and a series of indicators of enlistment waivers.

Analysis methods

Analysis was carried out remotely by Harvard Medical School analysts on the secure University of Michigan

Army STARRS Data Coordination Center server. Given that respondents differed in number of months of follow-up, we calculated survival curves for each outcome based on observed outcome distributions to estimate number of respondents who would have each outcome if they were all followed 24 months after enlistment (24-month morbid risk) using the actuarial method (Halli & Rao, 1992) implemented in SAS Proc LIFETEST (SAS Institute Inc., 2010). We projected morbid risk to 24 months despite our data going to 33 months because the number of soldiers followed beyond 24 months was too small for projection.

Discrete-time survival analysis with person-month the unit of analysis and a logistic link function (Willett & Singer, 1993) was used to develop a separate prediction model for first occurrence of each outcome. As noted above, our goal was to maximize classification rather than to test hypotheses about specific predictors, leading us to consider all potential predictors in the models. The major danger in doing this was overfitting (Ritchie, 2005; Upstill-Goddard *et al.* 2013). We addressed this problem by using the elastic net penalized regression method (Zou & Hastie, 2005) implemented in the R-package *glmnet* (Friedman *et al.* 2010) to select an optimal subset of predictors for each final model. Penalized regression methods are designed either to use shrinkage to include multiple highly correlated predictors in a single model, to select the most stable single predictor in each highly correlated set to represent all predictors in the set, or to use some combination of both approaches in the service of maximizing out-of-sample classification accuracy at the expense of coefficient accuracy. Given that performance-based neurocognitive test data collection was time-consuming (20 min administration time) and required special software, elastic net was implemented both with and without the neurocognitive measures to evaluate their incremental importance.

Prior to using elastic net, univariate associations of each potential predictor were estimated with each outcome controlling BCT site and time of data collection in SAS *proc logistic* (SAS Institute Inc., 2010). Functional forms of significant non-dichotomous predictors were transformed to capture simple nonlinearities. The elastic net analysis was then limited to significant univariate predictors. Once final elastic net models were estimated, individual-level predicted probabilities were calculated for each outcome and area under the receiver operating characteristic curve (AUC) was generated from these predicted probabilities to evaluate overall model fit. The sample was then divided into 20 groups of equal size (ventiles) for each outcome based on predicted probabilities of the outcome. When concentration of risk (COR; the observed proportion of realizations of a given outcome in a given

Table 1. 24-month morbid risk per 1000 soldiers and incidence per 1000 person-years of adverse outcomes in the New Soldiers Study ($n = 21832$)^a

	24-month morbid risk/1000 soldiers ^b	Incidence/1000 person-years ^c	S.E.	(<i>n</i>)
I. Mental–physical disorders				
Suicide attempt	8.2	5.1	0.4	(169)
Mental hospitalization	34.0	21.8	0.8	(739)
Positive drug test	18.9	11.9	0.7	(407)
Traumatic brain injury	2.6	1.8	0.3	(62)
Other severe injury	29.9	19.1	0.9	(658)
II. Violence				
Major physical perpetration (men)	3.8	2.5	0.3	(71)
Minor violence perpetration	11.2	7.0	0.5	(234)
Sexual assault perpetration (men)	5.2	3.1	0.4	(88)
Minor violence victimization	7.3	4.6	0.4	(162)
Sexual assault victimization (women)	37.7	25.1	2.6	(118)
III. Army career				
Attrition	189.8	119.0	2.9	(4285)
Demotion	60.8	39.2	1.2	(1337)

S.E., standard error; *n*, number of observed cases of the outcome in the sample.

^a Although the same sample of new soldiers was used for all outcomes, the number of person-months varied across outcomes because we predicted first occurrence of each outcome and each sample was censored separately either after the month when the outcome first occurred, termination of Regular Army service, or December 2013, whichever came first. The range of person-months was between 52 842 (to predict sexual assault victimization among women) and 4 20 706 (to predict attrition).

^b Morbid risk was estimated as the number of new soldiers predicted to have each outcome within 24 months of the beginning of service based on an actuarial projection (Halli & Rao, 1992). The projection was made to 24 months even though we had data for up to 33 months for some respondents because the number of respondents became too small for estimation beyond 24 months.

^c Incidence was estimated from the person-month file as the observed proportion of person-months with a realization of the outcome multiplied by 12 000 (month-to-year conversion \times 1000). There is no necessary relationship between incidence and morbid risk, as the former is based on a dataset that extends for between 13 and 33 months and depends on the timing of occurrences, whereas the former is projected only to 24 months and would have the same value whether outcomes occurred early or late in that time period.

ventile) was at least 15% among soldiers in the top-risk ventile (i.e. three times the expected value), we examined model coefficients and inspected COR across all 20 ventiles of predicted risk of that outcome.

Results

Morbid risk and correlations among outcomes

The most common outcome was attrition, with a 24-month morbid risk of 189.8/1000 soldiers (Table 1). Demotion had the next highest morbid risk (60.8/1000 soldiers), followed by sexual assault victimization (37.7/1000 female soldiers), mental hospitalization (34.0/1000 soldiers), severe injury (29.9/1000 soldiers), positive drug test (18.9/1000 soldiers), and minor violence perpetration (11.2/1000 soldiers). All other outcomes had 24-month morbid risk <10.0/1000 soldiers. Several outcomes were strongly inter-correlated (Table 2), with tetrachoric correlations of 0.84 for suicide attempt with mental hospitalization, 0.48–0.76 for major

physical perpetration with the other two perpetration outcomes, 0.54 for minor violence perpetration with minor violence victimization, and 0.55–0.70 for positive drug test with attrition and demotion.

Model accuracy

The number of predictors selected by elastic net was 3–29 (median = 14) across outcomes. AUC for the models ranged between 0.62 (mental hospitalization, severe injury) and 0.80 (TBI) (Table 3). Focusing on the nine outcomes with top-ventile COR above the minimum pre-specified level of 15%, 32.1–38.2% of new soldiers with three outcomes (major physical and sexual assault perpetration, TBI) and 21.6–29.8% of those with four other outcomes (minor violence perpetration, sexual assault victimization, suicide attempt, positive drug test) were among the 5% in the highest predicted risk ventiles for those outcomes (Fig. 1). The 40.5–46.5% of new soldiers with each of four outcomes (major physical and sexual assault perpetration, suicide

Table 2. Tetrachoric correlation matrix for all 12 observed outcomes in the total sample ($n = 21\,832$)^a

	a	b	c	d	e	f	g	h	i	j	k	l
I. Violence												
a. Major physical perpetration (men)	–											
b. Minor violence perpetration	0.76	–										
c. Sexual assault perpetration (men)	0.48	0.26	–									
d. Minor violence victimization	0.41	0.54	0.15	–								
e. Sexual assault victimization (women)	– ^b	–0.15	– ^b	0.48	–							
II. Mental-physical health												
f. Traumatic brain injury	0.08	–0.05	0.06	0.07	0.06	–						
g. Mental hospitalization	0.29	0.27	0.15	0.20	0.18	0.24	–					
h. Suicide attempt	0.27	0.23	0.17	0.15	0.21	0.10	0.84	–				
i. Positive drug test	0.36	0.18	0.16	0.17	0.23	0.16	0.31	0.22	–			
j. Severe injury	–0.01	0.00	0.08	–0.14	–0.01	0.18	0.18	0.17	–0.05	–		
III. Army career												
k. Attrition	0.21	0.16	0.15	–0.10	0.24	–0.09	0.40	0.42	0.55	–0.07	–	
l. Demotion	0.34	0.40	0.33	0.20	0.22	0.17	0.29	0.23	0.70	0.07	0.39	–

^a Correlations are estimated at the person-level ignoring duration of the follow-up period.

^b These correlations could not be estimated because these involve opposite-sex sex-specific outcomes.

Table 3. Performance of the final discrete-time survival model for each outcome^a

	AUC	Concentration of risk in ventiles predicted to have highest risk ^b			24-month morbid risk/1000 soldiers in ventiles predicted to have highest risk ^c		
		Top 5%	Top 10%	Top 15%	Top 5%	Top 10%	Top 15%
I. Mental-physical disorders							
Suicide attempt	0.74	29.8	40.5	48.9	48.9	33.2	26.7
Mental hospitalization	0.62	15.2	23.8	30.0	103.4	80.9	68.0
Positive drug test	0.71	21.6	29.7	41.4	81.6	56.1	52.2
Traumatic brain injury	0.80	38.2	46.5	55.8	19.9	12.1	9.7
Other severe injury	0.62	11.2	34.4	34.4	67.0	102.9	68.6
II. Violence							
Major physical perpetration (men)	0.78	34.0	45.8	47.7	25.8	17.4	12.1
Minor violence perpetration	0.76	24.0	35.9	45.8	53.8	40.2	34.3
Sexual assault perpetration (men)	0.78	32.1	44.6	57.2	33.4	23.2	19.8
Minor violence victimization	0.68	16.5	26.0	38.1	24.1	19.0	18.5
Sexual assault victimization (women)	0.71	23.1	35.5	40.3	174.2	133.8	101.3
III. Army career							
Attrition	0.65	13.2	21.6	28.4	501.1	410.0	359.4
Demotion	0.65	11.2	19.8	27.6	136.2	120.4	111.9

AUC, area under the receiver operating characteristic curve.

^a Each ventile represents 5% of the soldiers in the sample ranked in terms of their predicted risk of each outcome

^b Concentration of risk refers to the percent of all observed occurrences of the outcome in a ventile or ventiles of the predicted risk distribution.

^c Morbid risk was defined as the number of new soldiers within a ventile of predicted risk subsequently predicted to have each outcome within 24 months of the beginning of service based on the actuarial method. The projection was made to 24 months even though we had data for up to 33 months for some respondents because the number of respondents became too small for estimation beyond 24 months.




Fig. 1. Proportion of outcome observations within each ventile of predicted risk derived from final model[†].

attempt, TBI) were among the 10% in the two highest ventiles for those outcomes; and 55.8–57.2% of new soldiers with two outcomes (sexual assault perpetration, TBI) were among the 15% in the three highest ventiles for those outcomes. Morbid risk among new soldiers in the highest-risk ventile ranged widely due to the variations in overall morbid risk and COR, from a high of 501.1/1000 soldiers for attrition to a low of 19.9/1000 for TBI.

Model predictors

As noted above in the section on analysis methods, penalized regression maximizes classification accuracy at the expense of coefficient accuracy, making it important to focus more on model performance than on the specific predictors that entered the models. It is nonetheless noteworthy that 40.7% of the 772 potential predictors had significant (0.05-level, two-sided test) univariate associations with mental hospitalization, followed by 25.9–26.2% with suicide attempt and positive drug test, 16.7–17.9% with minor violence perpetration and sexual assault victimization, and no more than chance (4.8–6.3%) with the other outcomes (Table 4). As the potential predictors were unequally distributed across classes, we focused on observed/expected (O/E) predictor ratios in final models. Personality measures were proportionally overrepresented in the final models for seven outcomes, sociodemographics for six, stressors and administrative variables for three, mental disorders for two, and the

other classes of predictors for 0–1 outcomes (odds ratios and 95% confidence intervals of predictors in final models are available on request).

Several predictors selected by elastic net emerged in two or more final models. The common sociodemographics associated with elevated risk included minority status (predicting all violence perpetration outcomes, TBI, positive drug test), female sex (minor violence victimization, suicide attempt), low education (minor violence perpetration, positive drug test), and high religiosity (sexual assault perpetration, positive drug test). The most important mental disorder predictors associated with increased outcome risk were anger attacks (predicting major physical and minor violence perpetration, positive drug test), substance use disorders (minor violence perpetration, mental hospitalization, positive drug test), insomnia (minor violence perpetration and victimization, positive drug test), childhood behavioral disorders (minor violence perpetration, sexual assault perpetration and victimization, TBI, positive drug test), anxiety disorders (sexual assault perpetration, TBI, mental hospitalization, suicide attempt), total number of lifetime disorders (major physical and minor violence perpetration and victimization, mental hospitalization, suicide attempt), and lifetime treatment of mental disorders (major physical perpetration, suicide attempt).

The most important stressors included various chronic strains that occurred in the year prior to enlistment (predicting major physical perpetration, minor violence victimization, TBI), childhood physical abuse and physical assault victimization (sexual assault perpetration, TBI, positive drug test), family history of mental illness (sexual assault perpetration and

[†] The notes appear after the main text.

Table 4. Proportion of significant univariate predictors and final-model predictors across the eight predictor categories^a

Outcome	Sociodemographic		Mental disorders		Stressors		Personality		Social networks		Self-harm		Neurocognitive		Administrative		Total	
	% ^U ^b	O/E ^c	% ^d	(n) ^e														
I. Mental or physical disorders																		
Suicide attempt	10.5	1.2	34.3	1.2	18.9	0.6	58.2	1.8	29.7	0.5	15.4	1.7	62.5	–	10.8	0.9	25.9	(23)
Mental hospitalization	13.2	–	61.4	0.9	33.3	1.0	70.9	2.6	42.2	–	28.2	5.4	25.0	–	8.1	–	40.7	(11)
Positive drug test	21.1	1.9	38.2	0.9	21.6	0.6	29.1	1.3	28.1	2.2	5.1	–	–	–	13.5	–	26.2	(22)
Traumatic brain injury	5.3	0.9	3.9	0.4	12.6	1.6	3.6	0.9	1.6	–	–	–	25.0	12.9	2.7	1.4	6.3	(15)
II. Violence																		
Major physical perpetration (men)	7.0	0.7	3.4	1.0	5.0	0.7	3.6	1.4	3.1	–	–	–	50.0	–	13.5	6.2	5.2	(10)
Minor violence perpetration	7.0	1.1	26.2	2.0	10.4	0.2	18.2	0.8	23.4	0.7	2.6	–	87.5	–	10.8	1.2	16.7	(18)
Sexual assault perpetration (men)	7.0	2.1	1.3	0.3	5.0	0.8	14.5	3.2	3.1	0.9	2.6	1.5	37.5	–	2.7	–	4.8	(13)
Minor violence victimization	1.8	1.2	5.6	0.9	5.4	1.3	10.9	2.6	4.7	–	2.6	–	25.0	–	–	–	5.1	(11)
Sexual assault victimization (women)	7.9	1.1	16.7	0.6	26.6	2.0	49.1	1.2	1.6	–	7.7	–	–	–	–	–	17.9	(12)
% of all potential predictors	14.8		30.2		28.8		7.1		8.3		5.1		1.0		4.8			
(n) potential predictors	(114)		(233)		(222)		(55)		(64)		(39)		(8)		(37)			(772)

Neurocog, neurocognitive; Admin, administrative Army career variables; U, univariate; O/E, observed number of final model predictors divided by expected number of final model predictors; FM, final model; P, proportion; N, number.

^a Results are shown for the nine outcomes that achieved a concentration of risk in the top-ventile of risk at least three times the expected value (i.e. 15.0% ≤; see Table 3). Empty cells (–) indicate predictor categories that had no significant univariate associations or predictor categories that were not represented in a final model. Coefficients (odds ratios) for the final model predictors are available on request.

^b These columns report the percent of predictors within each predictor category with significant univariate associations with outcome (0.05 level, two-sided tests).

^c This ratio refers to the observed representation of the column predictor category divided by the expected representation of the column predictor category. For example, there was one sociodemographic variable selected among the 10 total final model variables for major physical perpetration – 10% representation. However, 114 of all predictors were sociodemographic variables. – 14.8%. This ratio (10.0/14.8) is equal to 0.7.

^d This column reports the total proportion of all predictors with significant univariate associations with each outcome.

^e This column reports the total number of final model predictors selected by elastic net.

victimization, TBI, mental hospitalization, suicide attempt), living in a foster home as a child (minor violence and sexual assault victimization), and number of lifetime stressors (all mental–physical disorder outcomes). Although a broad range of personality variables entered the different final models, only two were included in 2+ models: extroverted personality (predicting elevated risk in all three perpetration models); and non-secure attachment styles (predicting elevated risk of sexual assault perpetration, minor violence victimization, TBI, mental hospitalization, suicide attempt).

The remaining four groups of predictors were much less prominent in the final models. The only consistent predictor involving social networks was information about number of sexual partners (predicting elevated risk of minor violence perpetration, sexual assault perpetration, and positive drug tests). Indicators of self-harm were associated with elevated risk of sexual assault perpetration, mental hospitalization, and suicide attempt. The only administrative variable entering the models for multiple outcomes was the enlistment AFQT score, which inversely predicted risk of major physical and minor violence perpetration, TBI, and suicide attempts. Only one neurocognitive measure entered any final model: composite test efficiency predicted increased risk of TBI and improved top-ventile COR by 8.0% (from 38.2% to 46.2%).

Cross-outcome risk

Not surprisingly given that some outcomes were significantly inter-correlated, some composite predicted risk scores based on the final models were also significantly inter-correlated (detailed results available on request), raising a question whether predicted risk scores of Outcome A ever incrementally predicted Outcome B over the predict risk score for Outcome B. We explored this possibility by estimating models where each outcome was regressed on the predicted risk score for that outcome in addition to the predicted risk scores for the other outcomes. Only a handful of cross-outcome predictions were statistically significant and none improved COR in high-risk ventiles (detailed results available on request).

Discussion

We found that self-report questionnaire data collected at the beginning of service can be used to develop risk models with high top-ventile concentrations of risk for a number of subsequent negative soldier outcomes in the early years of service. Of particular note: 40.5% of suicide attempts occurred among the 10% of new soldiers with highest predicted risk of

that outcome, 34.0% of male major physical perpetrators were among the 5% with highest predicted risk of that outcome, 57.2% of male sexual assault perpetrators were among the 15% with highest predicted risk of that outcome, and 35.5% of female sexual assault victims were among the 10% with highest predicted risk of that outcome.

Caution is needed in interpreting the importance of the specific predictors in our final models because penalized regression methods maximize overall model performance at the expense of individual coefficient accuracy. Nonetheless, three observations are noteworthy about these predictors.

First, the vast majority of the signs of the associations between predictors and outcomes are consistent with those in prior studies of military personnel (Suris & Lind, 2008; Elbogen *et al.* 2010; Turchik & Wilson, 2010; Nock *et al.* 2013; Afifi *et al.* 2016) and civilians (Dahlberg, 1998; Kilpatrick *et al.* 2000; Kirst *et al.* 2014; Ryttila-Manninen *et al.* 2014). The major exception is that religiosity is positively associated with risk of sexual assault perpetration and positive drug tests. Religiosity is usually (Miller *et al.* 2000; Nonnemaker *et al.* 2003; Salas-Wright *et al.* 2012) found to be protective against these outcomes. However, at least three previous studies found religiosity to be associated with increased risk of similar outcomes (Jeffords, 1984; Schensul & Burkholder, 2005; Herman-Stahl *et al.* 2007). An understanding of why this might be the case would require more focused investigation.

Second, personality is the only predictor category with an O/E ratio consistently greater than 1.0 across outcomes. Few previous studies have examined personality predictors of negative soldier outcomes (MacManus *et al.* 2012a, b). Given our results, though, additional research might be warranted to compare predictive validity of the personality measures in the NSS with personality measures collected independently by the Army (Drasgow *et al.* 2012) as pre-accession screens for negative outcomes (Niebuhr *et al.* 2013).

Third, despite some evidence that objective measures of psychological characteristics have incremental validity over self-reports (Fuentes *et al.* 2006; Back *et al.* 2009; Huntjens *et al.* 2014), the performance-based neurocognitive test variables considered here did not figure prominently in our final models despite having high proportions of significant univariate associations. This raises a question about the value of including these tests in future surveys of new soldiers.

Three study limitations are noteworthy. First, the NSS was described to new soldiers as an independent survey in which individual-level responses would not be shared with Army leaders. If model results of the sort reported here are used to target preventive

interventions in future cohorts of new soldiers, results would have to be shared with Army leaders and respondents would have to be made aware of this fact before participating. This shift in auspices might alter reports in ways that reduce model performance. The STARRS results are nonetheless valuable in showing that respondent self-reports unencumbered by concerns about disclosure are capable of predicting the outcomes considered here with good accuracy. Based on these results, future studies might experiment with different approaches to motivate honest reporting in the context of respondents being told that results will be used to select soldiers for preventive interventions.

Second, our model results might not generalize beyond the 13–33 months of follow-up considered here and perhaps not beyond the 24 months for which we have a substantial sample. Replication of the current analyses over longer follow-up periods is needed to investigate this issue. Importantly, administrative data become richer over time, making it of interest in future long-term studies to examine the joint associations of baseline self-report measures and ongoing administrative measures in predicting the outcomes considered here.

Third, our use of administratively-recorded outcomes means that we excluded outcomes not reported to authorities (e.g. unreported sexual assault victimizations) and not detected by authorities (e.g. crime perpetrators who eluded authorities). Concern about this limitation is dampened, though, by two considerations. One is that administratively-recorded cases are often more severe than unrecorded cases. This is true, for example, of suicide attempts, where 100% of the severe attempts (i.e. those requiring hospitalization) need medical attention and are recorded in administrative records, even though some unknown number of less severe attempts are not known to authorities. The other is that administratively-recorded prevalence of these outcomes is high enough that prevention only of those cases would be of considerable value. Such interventions would also be expected to prevent some unreported cases, leading to conservative estimates of intervention cost-effectiveness based on administrative outcomes alone.

Conclusions

Within the context of these limitations, our results show that small subsets of new soldiers are responsible for high proportions of many of the negative outcomes considered here and that these high-risk new soldiers could be pinpointed at the beginning of their Army careers with models based on self-report data and basic administrative variables. These results argue strongly for the potential value of using self-report

surveys with new soldiers to target preventive interventions if the issue of confidentiality could be addressed successfully. Of course, the ultimate value of prediction models based on such surveys depends on the broadly-defined costs (both direct costs and competing risks) of these outcomes to the Army and the individual soldier and the effectiveness of such interventions in terms of number needed to treat (NNT; i.e., the number of high-risk new soldiers who would need to be treated to prevent one instance of a focal outcome). However, as noted in the introduction, a number of interventions exist that are very promising both in terms of costs and effectiveness when considered for targeted implementation, making the results reported here of considerable value.

Note

- ¹ Ventiles are 20 groups created by dividing the total sample into equally sized groups defined by rank order of predicted risk from the final models. Only the nine final models that had a concentration of risk in the top-ventile of risk at least three times the expected value are presented here.

Supplementary material

The supplementary material for this article can be found at <https://doi.org/10.1017/S003329171700071X>.

Acknowledgements

Army STARRS was sponsored by the Department of the Army and funded under cooperative agreement (R.J.U. & M.B.S., cooperative agreement number U01MH087981) (2009–2015) with the U.S. Department of Health and Human Services, National Institutes of Health, National Institute of Mental Health (NIH/NIMH); Subsequently, STARRS-LS was sponsored and funded by the Department of Defense (R.J.U. & M.B.S., USUHS grant number HU0001-15-2-0004); This research was also funded by the Department of Defense, Office of the Assistant Secretary for Defense for Health Affairs, Defense Health Program (OASD/HA), awarded and administered by the U.S. Army Medical Research & Materiel Command (USAMRMC), at Fort Detrick, MD (R.C.K., under contract numbers W81XWH-12-2-0113, W81XWH-13-1-0037); and Dr Street's effort was supported with resources and the use of facilities at VA Boston Healthcare System, Boston, MA.

Declaration of Interest

Dr Stein reports being paid as a consultant by Actelion Pharmaceuticals, Dart Neuroscience, Janssen

Pharmaceuticals, Resilience Therapeutics, and Oxeia Biopharmaceuticals; and for editorial work for UpToDate and the journal *Biological Psychiatry*. Dr Monahan is co-owner of Classification of Violence Risk (COVR), Inc. In the past 3 years, Dr Kessler received support for his epidemiological studies from Sanofi Aventis; was a consultant for Johnson & Johnson Wellness and Prevention, Shire, Takeda; and served on an advisory board for the Johnson & Johnson Services Inc. Lake Nona Life Project. Kessler is a co-owner of DataStat, Inc., a market research firm that carries out healthcare research. The remaining authors declare no conflict of interest.

Ethical Standards

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

Group Information

The Army STARRS Team consists of Co-Principal Investigators: Robert J. Ursano, MD (Uniformed Services University of the Health Sciences) and Murray B. Stein, MD, MPH (University of California San Diego and VA San Diego Healthcare System). Site Principal Investigators: Steven Heeringa, Ph.D. (University of Michigan) and Ronald C. Kessler, Ph.D. (Harvard Medical School). National Institute of Mental Health (NIMH) collaborating scientists: Lisa J. Colpe, Ph.D., MPH and Michael Schoenbaum, Ph.D. Army liaisons/consultants: COL Steven Cersovsky, MD, MPH [USAPHC (Provisional)] and Kenneth Cox, MD, MPH [USAPHC (Provisional)]. Other team members: Pablo A. Aliaga, MA (Uniformed Services University of the Health Sciences); COL David M. Benedek, MD (Uniformed Services University of the Health Sciences); K. Nikki Benevides, MA (Uniformed Services University of the Health Sciences); Paul D. Bliese, Ph.D. (University of South Carolina); Susan Borja, Ph.D. (NIMH); Evelyn J. Bromet, Ph.D. (Stony Brook University School of Medicine); Gregory G. Brown, Ph.D. (University of California San Diego); Laura Campbell-Sills, Ph.D. (University of California San Diego); Catherine L. Dempsey, Ph.D., MPH (Uniformed Services University of the Health Sciences); Carol S. Fullerton, Ph.D. (Uniformed Services University of the Health Sciences); Nancy Gebler, MA (University of Michigan); Robert K. Gifford, Ph.D. (Uniformed Services University of the Health Sciences); Stephen E. Gilman, ScD (Harvard School of Public Health); Marjan G. Holloway, Ph.D. (Uniformed Services University of the Health Sciences); Paul

E. Hurwitz, MPH (Uniformed Services University of the Health Sciences); Sonia Jain, Ph.D. (University of California San Diego); Tzu-Cheg Kao, Ph.D. (Uniformed Services University of the Health Sciences); Karestan C. Koenen, Ph.D. (Columbia University); Lisa Lewandowski-Romps, Ph.D. (University of Michigan); Holly Herberman Mash, Ph.D. (Uniformed Services University of the Health Sciences); James E. McCarroll, Ph.D., MPH (Uniformed Services University of the Health Sciences); James A. Naifeh, Ph.D. (Uniformed Services University of the Health Sciences); Tsz Hin Hinz Ng, MPH (Uniformed Services University of the Health Sciences); Matthew K. Nock, Ph.D. (Harvard University); Rema Raman, Ph.D. (University of California San Diego); Holly J. Ramsawh, Ph.D. (Uniformed Services University of the Health Sciences); Anthony Joseph Rosellini, Ph.D. (Harvard Medical School); Nancy A. Sampson, BA (Harvard Medical School); CDR Patcho Santiago, MD, MPH (Uniformed Services University of the Health Sciences); Michaelle Scanlon, MBA (NIMH); Jordan W. Smoller, MD, ScD (Harvard Medical School); Amy Street, Ph.D. (Boston University School of Medicine); Michael L. Thomas, Ph.D. (University of California San Diego); Leming Wang, MS (Uniformed Services University of the Health Sciences); Christina L. Wassel, Ph.D. (University of Vermont); Simon Wessely, FMedSci (King's College London); Christina L. Wryter, BA (Uniformed Services University of the Health Sciences); Hongyan Wu, MPH (Uniformed Services University of the Health Sciences); LTC Gary H. Wynn, MD (Uniformed Services University of the Health Sciences); and Alan M. Zaslavsky, Ph.D. (Harvard Medical School).

Disclaimer

The contents are solely the responsibility of the authors and do not necessarily represent the views of the Department of Health and Human Services, NIMH, or the Department of the Army, or the Department of Defense. No official endorsement should be made.

References

- Afifi TO, Taillieu T, Zamorski MA, Turner S, Cheung K, Sareen J (2016). Association of child abuse exposure with suicidal ideation, suicide plans, and suicide attempts in military personnel and the general population in Canada. *JAMA Psychiatry* 73, 229–238.
- Back MD, Schmukle SC, Egloff B (2009). Predicting actual behavior from the explicit and implicit self-concept of personality. *Journal of Personality and Social Psychology* 97, 533–548.
- Booth-Kewley S, Highfill-McRoy RM, Larson GE, Garland CF (2010). Psychosocial predictors of military misconduct. *Journal of Nervous and Mental Disease* 198, 91–98.

- Bulzacchelli MT, Sulsky SI, Rodriguez-Monguio R, Karlsson LH, Hill MO** (2014). Injury during U.S. Army basic combat training: a systematic review of risk factor studies. *American Journal of Preventive Medicine* **47**, 813–822.
- Canham-Chervak M, Hooper TI, Brennan Jr. FH, Craig SC, Girasek DC, Schaefer RA, Barbour G, Yew KS, Jones BH** (2010). A systematic process to prioritize prevention activities sustaining progress toward the reduction of military injuries. *American Journal of Preventive Medicine* **38** (1 Suppl.), S11–S18.
- Cassidy JD, Carroll LJ, Peloso PM, Borg J, von Holst H, Holm L, Kraus J, Coronado VG** (2004). Incidence, risk factors and prevention of mild traumatic brain injury: results of the WHO Collaborating Centre Task Force on Mild Traumatic Brain Injury. *Journal of Rehabilitation Medicine* **36** (43 Suppl.), 28–60.
- Dahlberg LL** (1998). Youth violence in the United States: major trends, risk factors, and prevention approaches. *American Journal of Preventive Medicine* **14**, 259–272.
- Department of Defense** (2014a). Department of Defense 2014–2016 Sexual Assault Prevention Strategy. Department of Defense: Arlington, Virginia (http://sapr.mil/public/docs/prevention/DoD_SAPR_Prevention_Strategy_2014-2016.pdf). Accessed 1 February 2016.
- Department of Defense** (2014b). DoD Workplace Violence Prevention and Response Policy. Department of Defense: Arlington, Virginia (<http://www.dtic.mil/whs/directives/corres/pdf/143806p.pdf>). Accessed 1 February 2016.
- Department of the Army** (2015a). AR-600-24 Health Promotion, Risk Reduction, and Suicide Prevention. Department of the Army: Washington, DC (http://www.lewis-mcchord.army.mil/dhr/asap/Doc/14%20APR%2015%20p600_24.pdf). Accessed 1 February 2016.
- Department of the Army** (2015b). AR-600-63 Army Health Promotion. Department of the Army: Washington, DC (https://www.army.mil/e2/downloads/rv7/r2/policydocs/r600_63.pdf). Accessed 1 February 2016.
- Department of the US Army** (2010). *Army Health Promotion, Risk Reduction, and Suicide Prevention: Report 2010*. US Army: Washington, DC.
- Department of the US Army** (2012). *Army 2020: Generating Health & Discipline in the Force ahead of the Strategic Reset*. US Army: Washington, DC.
- Dragow F, Stark S, Chernyshenko OS, Nye CD, Hulin CL, White LA** (2012). Technical Report 1311 – Development of the Tailored Adaptive Personality Assessment System (TAPAS) to Support Army Selection and Classification Decisions. U.S. Army Research Institute for the Behavioral and Social Sciences: Fort Belvoir, Virginia (<http://www.dtic.mil/cgi-bin/GetTRDoc?AD=ADA564422>). Accessed 1 February 2016.
- Elbogen EB, Fuller S, Johnson SC, Brooks S, Kinneer P, Calhoun PS, Beckham JC** (2010). Improving risk assessment of violence among military veterans: an evidence-based approach for clinical decision-making. *Clinical Psychology Review* **30**, 595–607.
- Elmasry H, Boivin MR, Feng X, Packnett ER, Cowan DN** (2017). Preenlistment and early service risk factors for traumatic brain injury in the Army and Marine Corps: FY 2002–2010. *Journal of Head Trauma Rehabilitation* **32**, E1–E7.
- Foster EM, Jones D** (2006). Can a costly intervention be cost-effective?: an analysis of violence prevention. *Archives of General Psychiatry* **63**, 1284–1291.
- Friedman J, Hastie T, Tibshirani R** (2010). Regularization paths for generalized linear models via coordinate descent. *Journal of Statistical Software* **33**, 1–22.
- Fuentes D, Tavares H, Artes R, Gorenstein C** (2006). Self-reported and neuropsychological measures of impulsivity in pathological gambling. *Journal of the International Neuropsychological Society: IJNS* **12**, 907–912.
- Golubnitschaja O, Costigliola V** (2012). General report & recommendations in predictive, preventive and personalised medicine 2012: white paper of the European Association for Predictive, Preventive and Personalised Medicine. *EPMA Journal* **3**, 14.
- Halli SS, Rao KV** (1992). *Advanced Techniques of Population Analysis*. Plenum Press: New York, NY.
- Herman-Stahl MA, Krebs CP, Kroutil LA, Heller DC** (2007). Risk and protective factors for methamphetamine use and nonmedical use of prescription stimulants among young adults aged 18 to 25. *Addictive Behaviors* **32**, 1003–1015.
- Huntjens RJ, Rijkeboer MM, Krakau A, de Jong PJ** (2014). Implicit versus explicit measures of self-concept of self-control and their differential predictive power for spontaneous trait-relevant behaviors. *Journal of Behavior Therapy and Experimental Psychiatry* **45**, 1–7.
- Institute of Medicine** (2010). *Returning Home from Iraq and Afghanistan: Preliminary Assessment of Readjustment Needs of Veterans, Service Members, and their Families*. The National Academies Press: Washington, DC.
- Iribarren C, Sidney S, Jacobs Jr. DR, Weisner C** (2000). Hospitalization for suicide attempt and completed suicide: epidemiological features in a managed care population. *Social Psychiatry and Psychiatric Epidemiology* **35**, 288–296.
- Jeffords CR** (1984). The impact of sex-role and religious attitudes upon forced marital intercourse norms. *Sex Roles* **11**, 543–552.
- Kapp L** (2013). Recruiting and Retention: An Overview of FY2011 and FY2012 Results for Active and Reserve Component Enlisted Personnel. Congressional Research Service: Washington, DC (<https://www.fas.org/sgp/crs/natsec/RL32965.pdf>). Accessed 1 February 2016.
- Kaufman KR, Brodine S, Shaffer R** (2000). Military training-related injuries: surveillance, research, and prevention. *American Journal of Preventive Medicine* **18**, 54–63.
- Kessler RC, Warner CH, Ivany C, Petukhova MV, Rose S, Bromet EJ, Brown III M, Cai T, Colpe LJ, Cox KL, Fullerton CS, Gilman SE, Gruber MJ, Heeringa SG, Lewandowski-Romps L, Li J, Millikan-Bell AM, Naifeh JA, Nock MK, Rosellini AJ, Sampson NA, Schoenbaum M, Stein MB, Wessely S, Zaslavsky AM, Ursano RJ** (2015). Predicting suicides after psychiatric hospitalization in US Army soldiers: the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS). *JAMA Psychiatry* **72**, 49–57.

- Kilpatrick DG, Acierno R, Saunders B, Resnick HS, Best CL, Schnurr PP (2000). Risk factors for adolescent substance abuse and dependence: data from a national sample. *Journal of Consulting and Clinical Psychology* **68**, 19–30.
- Kirst M, Mecredy G, Borland T, Chaiton M (2014). Predictors of substance use among young adults transitioning away from high school: a narrative review. *Substance Use and Misuse* **49**, 1795–1807.
- Knapik JJ, Jones BH, Hauret K, Darakjy S, Piskator E (2004). A Review of the Literature on Attrition from the Military Services: Risk Factors for Attrition and Strategies to Reduce Attrition: USACHPPM REPORT NO. 12-HF-01Q3A-04. U.S. Army Center for Health Promotion and Preventative Medicine: Aberdeen Proving Ground, MD (<http://www.dtic.mil/dtic/tr/fulltext/u2/a427744.pdf>). Accessed 1 February 2016.
- Kubisiak UC, Lentz E, Horgen KE, Bryant RH, Connell PW, Tuttle MD, Borman WC, Young MC, Morath R (2009). ARI Research Note 2009–13. Review of Interventions for Reducing Enlisted Attrition in the U.S. Military: an Update. United States Army Research Institute for the Behavioral and Social Sciences: Arlington, VA (<http://www.dtic.mil/dtic/tr/fulltext/u2/a508188.pdf>). Accessed 1 February 2016.
- MacManus D, Dean K, Al Bakir M, Iversen AC, Hull L, Fahy T, Wessely S, Fear NT (2012a). Violent behaviour in U.K. military personnel returning home after deployment. *Psychological Medicine* **42**, 1663–1673.
- MacManus D, Dean K, Iversen AC, Hull L, Jones N, Fahy T, Wessely S, Fear NT (2012b). Impact of pre-enlistment antisocial behaviour on behavioural outcomes among UK military personnel. *Social Psychiatry and Psychiatric Epidemiology* **47**, 1353–1358.
- Miller L, Davies M, Greenwald S (2000). Religiosity and substance use and abuse among adolescents in the National Comorbidity Survey. *Journal of the American Academy of Child and Adolescent Psychiatry* **39**, 1190–1197.
- Moore TM, Gur RC, Thomas ML, Brown GG, Nock MK, Savitt AP, Keilp JG, Heeringa S, Ursano RJ, Stein MB (2017). Development, administration, and structural validity of a brief, computerized neurocognitive battery. *Assessment*. Published online 30 January 2017. doi: <http://dx.doi.org/10.1177%2F1073191116689820>.
- Niebuhr DW, Gubata ME, Oetting AA, Weber NS, Feng X, Cowan DN (2013). Personality assessment questionnaire as a pre-accession screen for risk of mental disorders and early attrition in U.S. Army recruits. *Psychological Services* **10**, 378–385.
- Nock MK, Deming CA, Fullerton CS, Gilman SE, Goldenberg M, Kessler RC, McCarroll JE, McLaughlin KA, Peterson C, Schoenbaum M, Stanley B, Ursano RJ (2013). Suicide among soldiers: a review of psychosocial risk and protective factors. *Psychiatry* **76**, 97–125.
- Nonnemaker J, McNeely C, Blum R (2003). Public and private domains of religiosity and adolescent health risk behaviors: evidence from the National Longitudinal Study of Adolescent Health. *Social Science and Medicine* **57**, 2049–2054.
- Parkkari J, Taanila H, Suni J, Mattila VM, Ohrankammen O, Vuorinen P, Kannus P, Pihlajamaki H (2011). Neuromuscular training with injury prevention counselling to decrease the risk of acute musculoskeletal injury in young men during military service: a population-based, randomised study. *BMC Medicine* **9**, 35.
- Ritchie MD (2005). Bioinformatics approaches for detecting gene-gene and gene-environment interactions in studies of human disease. *Neurosurgical Focus* **19**, 1–4.
- Rosellini AJ, Heeringa SG, Stein MB, Ursano RJ, Chiu WT, Colpe LJ, Fullerton CS, Gilman SE, Hwang I, Naifeh JA, Nock MK, Petukhova M, Sampson NA, Schoenbaum M, Zaslavsky AM, Kessler RC (2015). Lifetime prevalence of DSM-IV mental disorders among new soldiers in the U.S. Army: results from the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS). *Depression and Anxiety* **32**, 13–24.
- Rosellini AJ, Monahan J, Street AE, Heeringa SG, Hill ED, Petukhova M, Reis BY, Sampson NA, Bliese P, Schoenbaum M, Stein MB, Ursano RJ, Kessler RC (2016). Predicting non-familial major physical violent crime perpetration in the US Army from administrative data. *Psychological Medicine* **46**, 303–316.
- Rudd MD, Bryan CJ, Wertenberger EG, Peterson AL, Young-McCaughan S, Mintz J, Williams SR, Arne KA, Breitbart J, Delano K, Wilkinson E, Bruce TO (2015). Brief cognitive-behavioral therapy effects on post-treatment suicide attempts in a military sample: results of a randomized clinical trial with 2-year follow-up. *American Journal of Psychiatry* **172**, 441–449.
- Rytila-Manninen M, Lindberg N, Haravuori H, Kettunen K, Marttunen M, Joukamaa M, Frojd S (2014). Adverse childhood experiences as risk factors for serious mental disorders and inpatient hospitalization among adolescents. *Child Abuse and Neglect* **38**, 2021–2032.
- Salas-Wright CP, Vaughn MG, Hodge DR, Perron BE (2012). Religiosity profiles of American youth in relation to substance use, violence, and delinquency. *Journal of Youth and Adolescence* **41**, 1560–1575.
- SAS Institute Inc. (2010). *SAS/STAT Software*. SAS Institute Inc.: Cary, NC.
- Schensul JJ, Burkholder GJ (2005). Vulnerability, social networks, sites, and selling as predictors of drug use among urban African American and Puerto Rican emerging adults. *Journal of Drug Issues* **35**, 379–408.
- Senn CY, Eliasziw M, Barata PC, Thurston WE, Newby-Clark IR, Radtke HL, Hobden KL (2015). Efficacy of a sexual assault resistance program for university women. *New England Journal of Medicine* **372**, 2326–2335.
- Shea MT, Lambert J, Reddy MK (2013). A randomized pilot study of anger treatment for Iraq and Afghanistan veterans. *Behaviour Research and Therapy* **51**, 607–613.
- Street AE, Rosellini AJ, Ursano RJ, Heeringa SG, Hill ED, Monahan J, Naifeh JA, Petukhova MV, Reis BY, Sampson NA, Bliese PD, Stein MB, Zaslavsky AM, Kessler RC (2016). Developing a risk model to target high-risk preventive interventions for sexual assault victimization among female US Army soldiers. *Clinical Psychological Science* **4**, 939–956.
- Suris A, Lind L (2008). Military sexual trauma: a review of prevalence and associated health consequences in veterans. *Trauma, Violence and Abuse* **9**, 250–269.
- Theodoroff SM, Lewis MS, Folmer RL, Henry JA, Carlson KF (2015). Hearing impairment and tinnitus: prevalence,

- risk factors, and outcomes in US service members and veterans deployed to the Iraq and Afghanistan wars. *Epidemiologic Reviews* 37, 71–85.
- Turchik JA, Wilson SM** (2010). Sexual assault in the U.S. military: A review of the literature and recommendations for the future. *Aggression and Violent Behavior* 15, 267–277.
- Upstill-Goddard R, Eccles D, Fliege J, Collins A** (2013). Machine learning approaches for the discovery of gene-gene interactions in disease data. *Briefings in Bioinformatics* 14, 251–260.
- Ursano RJ, Colpe LJ, Heeringa SG, Kessler RC, Schoenbaum M, Stein MB** (2014). The army study to assess risk and resilience in servicemembers (Army STARRS). *Psychiatry* 77, 107–119.
- U.S. Department of Justice** (2011). National Corrections Reporting Program, 2009 (ICPSR 30799). National Archive of Criminal Justice Data: Ann Arbor, MI (<http://www.icpsr.umich.edu/icpsrweb/NACJD/studies/30799?archive=NACJD&permit%5B0%5D=AVAILABLE&q=30799&x=0&y=0>). Accessed 1 February 2016.
- Vungkhanching M, Heinemann AW, Langley MJ, Ridgely M, Kramer KM** (2007). Feasibility of a skills-based substance abuse prevention program following traumatic brain injury. *Journal of Head Trauma Rehabilitation* 22, 167–176.
- Willett JB, Singer JD** (1993). Investigating onset, cessation, relapse, and recovery: why you should, and how you can, use discrete-time survival analysis to examine event occurrence. *Journal of Consulting Clinical Psychology* 61, 952–965.
- Zou H, Hastie T** (2005). Regularization and variable selection via the elastic net. *Journal of the Royal Statistical Society: Series B* 67, 301–320.